

BENWAY SCHOOL

2017-2018 SCHOOL YEAR

6.

OVER-THE-COUNTER MEDICATION FORM TO BE SIGNED BY A PHYSICIAN

One of the emergency/permission forms you signed was the MEDICATION CONSENT FORM. This form referred to ANY prescription medication and/or over-the-counter medication for your child. If your child requires a prescribed medication, daily or on a short term basis, we must have Medication Consent completed and signed by your physician.

However, from time-to-time your child may come to the office in need of relief for a cough, sore throat, indigestion, upset stomach, cramps, headache, or body ache. For that reason, we have listed below the **ONLY** over-the-counter medication we will supply **only if** your physician checks off a medication listed and signs below. **Please note that if this form is NOT signed and returned to the office, your child will not be administered any of the over-the-counter medications listed below.**

_____ Cough Drops

_____ Ibuprofen 200 MG tablets (also known as Advil or Motrin). Children 12 years or older take (swallow) 1 - 2 tablets for headache, cramps, or body aches. Do not exceed 6 tablets in 24 hours.

_____ Ibuprofen Junior strength 100 MG chewable tablets. Use dosing chart ages 6-8 (48-59lbs.) take 2 tablets; ages 9-10 (60-71 lbs) take 2.5 tablets; a 11 (72-95 lbs.) take (chew) 3 tablets. For headache, cramps or body aches. Repeat dose every 6-8 hrs. if symptoms persist. Do not exceed more than 4 doses in 24 hours.

_____ Acetaminophen pain reliever 500 MG gelcaps (also known as Tylenol). Adults and children 12 years or older take/swallow 2 gelcaps every 6 hours while symptoms like fever or body aches persist. Do not exceed 6 gelcaps in 24 hours.

_____ Acetaminophen 325 MG. Adults and children 12 years or older take 2-3 tablets every 6 hours as needed for headache or fever.

_____ Acetaminophen Junior Strength (Chewable). See weight and age chart for children under 12. For headaches or fever.

_____ Antacids (also known as Tums or Rolaids). Children 12 years and older take 1-2 tabs for acid reflux.

_____ Bismuth (also known as Pepto-Bismol). Children 12 years and older take 1-2 tablets (or 1 table spoon). For nausea, indigestion, diarrhea.

Student Name: _____ Parent/Guardian Signature: _____

Physician Name (Printed) _____ Physician Signature: _____

Physician Address & Phone Number or Stamp: _____

